

A new and elementary proof of Newton's "favorite" quadrature formulae

Cezar Lupu, Tudorel Lupu

In this note we shall give a new proof to a quadrature formulae due to Newton(see [1.]) which states

Theorem 0.1 *If $f : [a, b] \rightarrow \mathbb{R}$ is four times differentiable on $[a, b]$ with $f^{(4)}$ continuous, then there is some $\xi \in [a, b]$ such that*

$$\int_a^b f(x)dx = \frac{b-a}{8} \left[f(a) + 3f\left(\frac{2a+b}{3}\right) + 3f\left(\frac{2b+a}{3}\right) + f(b) \right] - \frac{(b-a)^5}{6480} f^{(4)}(\xi).$$

In order to prove this theorem we need to show first three lemmas starting with

Lemma 1. *If $f : [a, b] \rightarrow \mathbb{R}$ is four times differentiable with $f^{(4)}$ continuous, then there is some $\xi_1 \in [a, b]$ such that*

$$\int_a^b f(x)dx = (b-a)f\left(\frac{a+b}{2}\right) + \frac{(b-a)^3}{24} f^{(2)}\left(\frac{a+b}{2}\right) + \frac{(b-a)^5}{1920} f^{(4)}(\xi_1).$$

Proof. We begin with the following integral

$$\int_a^b (x-a)^4 \left[f^{(4)}\left(\frac{a+x}{2}\right) + f^{(4)}\left(\frac{2b+a-x}{2}\right) \right] dx.$$

Integrating by part we get

$$\begin{aligned} \int_a^b (x-a)^4 \left[f^{(4)}\left(\frac{a+x}{2}\right) + f^{(4)}\left(\frac{2b+a-x}{2}\right) \right] dx &= 2(x-a)^4 \left[f^{(3)}\left(\frac{a+x}{2}\right) - f^{(3)}\left(\frac{2b+a-x}{2}\right) \right] \Big|_a^b \\ &- 2 \int_a^b 4(x-a)^3 \left[f^{(3)}\left(\frac{a+x}{2}\right) - f^{(3)}\left(\frac{2b+a-x}{2}\right) \right] dx = -16(x-a)^3 \left[f^{(2)}\left(\frac{a+x}{2}\right) + f^{(2)}\left(\frac{2b+a-x}{2}\right) \right] \Big|_a^b \\ &+ 48 \int_a^b (x-a)^2 \left[f^{(2)}\left(\frac{a+x}{2}\right) + f^{(2)}\left(\frac{2b+a-x}{2}\right) \right] dx = -32(b-a)^3 f^{(2)}\left(\frac{a+b}{2}\right) + \\ &+ 96(x-a)^2 \left[f'\left(\frac{a+x}{2}\right) - f'\left(\frac{2b+a-x}{2}\right) \right] \Big|_a^b - 192 \int_a^b (x-a) \left[f'\left(\frac{a+x}{2}\right) - f'\left(\frac{2b+a-x}{2}\right) \right] dx \\ &= -32(b-a)^3 f^{(2)}\left(\frac{a+b}{2}\right) - 384(x-a) \left[f\left(\frac{a+x}{2}\right) + f\left(\frac{2b+a-x}{2}\right) \right] \Big|_a^b \\ &+ 384 \int_a^b \left[f\left(\frac{a+x}{2}\right) + f\left(\frac{2b+a-x}{2}\right) \right] dx = 768 \int_a^b f(x)dx - 768(b-a)f\left(\frac{a+b}{2}\right) - 32(b-a)^3 f^{(2)}\left(\frac{a+b}{2}\right). \end{aligned}$$

On the other hand, by the mean value theorem, there is $\eta \in [a, b]$ such that

$$\int_a^b (x-a)^4 \left[f^{(4)}\left(\frac{a+x}{2}\right) + f^{(4)}\left(\frac{2b+a-x}{2}\right) \right] dx = \left[f^{(4)}\left(\frac{a+\eta}{2}\right) + f^{(4)}\left(\frac{2b+a-\eta}{2}\right) \right] \int_a^b (x-a)^4 dx.$$

But, $f^{(4)}$ is continuous, so $f^{(4)}$ has intermediate value property. It follows that there is $\xi_1 \in [a, b]$ such that

$$f^{(4)}\left(\frac{a+\eta}{2}\right) + f^{(4)}\left(\frac{2b+a-\eta}{2}\right) = 2f^{(4)}(\xi_1),$$

so

$$\int_a^b (x-a)^4 \left[f^{(4)}\left(\frac{a+x}{2}\right) + f^{(4)}\left(\frac{2b+a-x}{2}\right) \right] dx = \frac{2(b-a)^5}{5} f^{(4)}(\xi_1).$$

We get

$$768 \int_a^b f(x) dx = 768(b-a)f\left(\frac{a+b}{2}\right) + 32(b-a)^3 f^{(2)}\left(\frac{a+b}{2}\right) + \frac{2(b-a)^5}{5} f^{(4)}(\xi_1)$$

which is equivalent to

$$\int_a^b f(x) dx = (b-a)f\left(\frac{a+b}{2}\right) + \frac{(b-a)^3}{24} f^{(2)}\left(\frac{a+b}{2}\right) + \frac{(b-a)^5}{1920} f^{(4)}(\xi_1),$$

where $\xi_1 \in [a, b]$. □

Lemma 2.

If $f : [a, b] \rightarrow \mathbb{R}$ is four times differentiable with $f^{(4)}$ continuous, then there is some $\xi_2 \in [a, b]$ such that

$$\int_a^b f(x) dx = \frac{(b-a)}{8} \left[f(a) + 6f\left(\frac{a+b}{2}\right) + f(b) \right] + \frac{(b-a)^3}{96} f^{(2)}\left(\frac{a+b}{2}\right) - \frac{(b-a)^5}{7680} f^{(4)}(\xi_2).$$

Proof. Like in the proof of lemma 1, we will begin with the following integral

$$\int_a^b (x-a)^3(x-b) \left[f^{(4)}\left(\frac{a+x}{2}\right) + f^{(4)}\left(\frac{2b+a-x}{2}\right) \right] dx.$$

Integrating by part we get

$$\begin{aligned} \int_a^b (x-a)^3(x-b) \left[f^{(4)}\left(\frac{a+x}{2}\right) + f^{(4)}\left(\frac{2b+a-x}{2}\right) \right] dx &= 2(x-a)^3(x-b) \left[f^{(3)}\left(\frac{a+x}{2}\right) - f^{(3)}\left(\frac{2b+a-x}{2}\right) \right] \Big|_a^b - \\ &- 2 \int_a^b [3(x-a)^2(x-b) + (x-a)^3] \left[f^{(3)}\left(\frac{a+x}{2}\right) - f^{(3)}\left(\frac{2b+a-x}{2}\right) \right] dx = \\ &= -4 [3(x-a)^2(x-b) + (x-a)^3] \left[f^{(2)}\left(\frac{a+x}{2}\right) + f^{(2)}\left(\frac{2b+a-x}{2}\right) \right] \Big|_a^b + \\ &+ 24 \int_a^b [(x-a)^2 + (x-a)(x-b)] \left[f^{(2)}\left(\frac{a+x}{2}\right) + f^{(2)}\left(\frac{2b+a-x}{2}\right) \right] dx = \\ &= -8(b-a)^3 f^{(2)}\left(\frac{a+b}{2}\right) + 48 [(x-a)^2 + (x-a)(x-b)] \left[f'\left(\frac{a+x}{2}\right) - f'\left(\frac{2b+a-x}{2}\right) \right] \Big|_a^b - \\ &- 48 \int_a^b [3(x-a) + x-b] \left[f'\left(\frac{a+x}{2}\right) - f'\left(\frac{2b+a-x}{2}\right) \right] dx = \\ &= -8(b-a)^3 f^{(2)}\left(\frac{a+b}{2}\right) - 96 [3(x-a) + (x-b)] \left[f\left(\frac{a+x}{2}\right) + f\left(\frac{2b+a-x}{2}\right) \right] \Big|_a^b \\ &+ 384 \int_a^b \left[f\left(\frac{a+x}{2}\right) + f\left(\frac{2b+a-x}{2}\right) \right] dx = 768 \int_a^b f(x) dx - 96 \cdot 6(b-a) f\left(\frac{a+b}{2}\right) \\ &- 96(b-a) [f(a) + f(b)] - 8(b-a)^3 f^{(2)}\left(\frac{a+b}{2}\right) = 768 \int_a^b f(x) dx - 96(b-a) \left[f(a) + 6f\left(\frac{a+b}{2}\right) + f(b) \right] - \end{aligned}$$

$-8(b-a)^3 f^{(2)}\left(\frac{a+b}{2}\right)$. On the other hand, by the mean value theorem there is some $\eta' \in [a, b]$ such that

$$\int_a^b (x-a)^3(x-b) \left[f^{(4)}\left(\frac{a+x}{2}\right) + f^{(4)}\left(\frac{2b+a-x}{2}\right) \right] dx = \left[f^{(4)}\left(\frac{a+\eta'}{2}\right) + f^{(4)}\left(\frac{2b+a-\eta'}{2}\right) \right] \cdot \int_a^b (x-a)^3(x-b) dx.$$

By the intermediate value property, there is some $\xi_2 \in [a, b]$ such that

$$f^{(4)}\left(\frac{a+\eta'}{2}\right) + f^{(4)}\left(\frac{2b+a-\eta'}{2}\right) = 2f^{(4)}(\xi_2).$$

A simple calculation shows that $\int_a^b (x-a)^3(x-b) dx = -\frac{(b-a)^5}{20}$. Now, combining these two equalities obtained, the lemma follows immediately. $\square$

Lemma 3(Simpson-Cavalieri).

If $f : [a, b] \rightarrow \mathbb{R}$ is four times differentiable with $f^{(4)}$ continuous, then there is some $\xi_3 \in [a, b]$ such that

$$\int_a^b f(x) dx = \frac{(b-a)}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{(b-a)^5}{2880} f^{(4)}(\xi_3).$$

Proof. This formulae can be proved in many ways, but we prefer to deduce it by eliminating $f^{(2)}\left(\frac{a+b}{2}\right)$ from lemma 1 and lemma 2. $\square$

Now, we shall prove the main result of our paper. First, we apply lemma 1 on the interval $\left[\frac{2a+b}{3}, \frac{2b+a}{3}\right]$. Then, there is some $\xi'_1 \in \left[\frac{2a+b}{3}, \frac{2b+a}{3}\right]$ such that

$$\int_{\frac{2a+b}{3}}^{\frac{2b+a}{3}} f(x) dx = \frac{b-a}{3} f\left(\frac{a+b}{2}\right) + \frac{(b-a)^3}{24 \cdot 27} f^{(2)}\left(\frac{a+b}{2}\right) + \frac{(b-a)^5}{1920 \cdot 3^5} f^{(4)}(\xi'_1). (*)$$

By eliminating $f^{(2)}\left(\frac{a+b}{2}\right)$ from lemma 2 and (*) we get

$$4 \int_a^b f(x) dx = 27 \int_{\frac{2a+b}{3}}^{\frac{2b+a}{3}} f(x) dx + \frac{b-a}{2} \left[f(a) - 12f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{(b-a)^5}{9 \cdot 192} f^{(4)}(\xi_4), (**)$$

with $\xi_4 \in [a, b]$. Now, applying Lemma 3(Simpson-Cavalieri), we get

$$\int_{\frac{2a+b}{3}}^{\frac{2b+a}{3}} f(x) dx = \frac{b-a}{18} \left[f\left(\frac{2a+b}{3}\right) + 4f\left(\frac{a+b}{2}\right) + f\left(\frac{2b+a}{3}\right) \right] - \frac{(b-a)^5}{3^5 \cdot 2880} f^{(4)}(\xi_5) (***)$$

with $\xi_5 \in \left[\frac{2a+b}{3}, \frac{2b+a}{3}\right]$. Now, replacing (***) in (**) we have

$$\begin{aligned} 4 \int_a^b f(x) dx &= \frac{27(b-a)}{18} \left[f\left(\frac{2a+b}{3}\right) + 4f\left(\frac{a+b}{2}\right) + f\left(\frac{2b+a}{3}\right) \right] - \frac{(b-a)^5}{9 \cdot 2880} f^{(4)}(\xi_5) \\ &\quad + \frac{(b-a)}{2} \left[f(a) - 12f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{(b-a)^5}{9 \cdot 192} f^{(4)}(\xi_4), \end{aligned}$$

with $\xi_4, \xi_5 \in [a, b]$. The above formulae is equivalent to

$$4 \int_a^b f(x) dx = \frac{b-a}{2} \left[f(a) + 3f\left(\frac{2a+b}{3}\right) + 3f\left(\frac{2b+a}{3}\right) + f(b) \right] - \left[\frac{(b-a)^5}{9 \cdot 2880} f^{(4)}(\xi_5) + \frac{(b-a)^5}{9 \cdot 192} f^{(4)}(\xi_4) \right].$$

Now, using intermediate value property, we get a $\xi \in [a, b]$ such that

$$\frac{(b-a)^5}{9 \cdot 2880} f^{(4)}(\xi_5) + \frac{(b-a)^5}{9 \cdot 192} f^{(4)}(\xi_4) = \frac{(b-a)^5}{1620} f^{(4)}(\xi).$$

Now, Newton's result follows immediately.

References.

- [1.] D.V. Ionescu-Restul în formula de cuadratură "preferată" a lui Newton, Gaz.Mat., vol 7, 1965, pag. 298-303.
- [2.] J.V. Steffensen-Interpolation, Baltimore, 1927
- [3.] B.P. Demidovich, I.A Maron-Computational Mathematics, Moscow: Mir Publishers, 1976. 2nd edition. Translated from Russian.

Cezar Lupu, student Faculty of Mathematics, University of Bucharest, Bucharest, Romania. e-mail: lupucezar@yahoo.com

Tudorel Lupu, teacher, Decebal High school, Constanza, Romania.